

# Methoden moderner Röntgenphysik I: Struktur und Dynamik kondensierter Materie

Vorlesung zum Haupt/Masterstudiengang Physik

WS 2008/9

G. Grübel, M. Martins, E. Weckert et al.

Location: SemRm 4, Physik, Jungiusstrasse

Thursdays 10.15 – 11.45

G.Grübel (GG), A.Meents (AM), C. Gutt (CG), S. Roth (SR)

# Methoden moderner Röntgenphysik I: Struktur und Dynamik kondensierter Materie

## Hard X-Rays – Kinematic Scattering Theory **Lecture 3**

23.10.	Introduction	(GG)
30.10.	Sources of X-rays, Refraction and Reflexion	(GG)
6.11.	Kinematic Scattering Theory	(GG)
13.11.	Small Angle and Anomalous Scattering	(GG)
20.11. - 11.12.	Modern Crystallography	(AM)
18.12. - 15. 1.	Coherence base techniques	(CG)
4.12. - 5. 2.	Soft Matter Applications	(SR)

# Coherence of light and matter I: from basic concepts to modern applications

Introduction into X-ray physics: 23.10.-13.11.

## Introduction

Overview, Introduction to X-ray Scattering (Scattering from atoms, crystals,..., absorption, reflection, coherence,...)

## Sources, Reflection and Refraction

Sources of X-rays, Refraction, reflection, Snell's law, Fresnel equations

## Kinematical Diffraction

Diffraction from an atom, molecule, crystal, reciprocal lattice, structure factor,..

## SAXS, Anomalous Diffraction


Introduction into small angle scattering and anomalous scattering

# Kinematical Diffraction

One of the main applications of X-rays is the determination of structure(s) using diffraction.

Assume the scattering to be weak; multiple scattering effects are to be neglected: weak scattering limit  $\equiv$  kinematical approximation.

Consider a 2 electron system:



$$z = r \cos \alpha; k z = k r \cos \alpha = \mathbf{k} \cdot \mathbf{r}$$

$$y = r \cos \beta; k'y = k' y \cos \beta = \mathbf{k}' \cdot \mathbf{r}$$

path- or phase difference:

$$\Delta \Phi = z - y = \mathbf{k} \cdot \mathbf{r} - \mathbf{k}' \cdot \mathbf{r} = \mathbf{Q} \cdot \mathbf{r}$$

with

$$Q = (4\pi/\lambda) \sin \theta$$

scattering amplitude for 2 electrons:

$$A(\mathbf{Q}) = -r_0 [1 + \exp(i\mathbf{Q}\mathbf{r})]$$

$$I(\mathbf{Q}) = A(\mathbf{Q}) A(\mathbf{Q})^*$$


$$= 2r_0^2 [1 + \cos(Qr)]$$

**see Fig. 4.2**

for many electrons:

$$A(\mathbf{Q}) = -r_0 \sum r_j \exp(i\mathbf{Q}\mathbf{r}_j)$$

# Kinematical Diffraction




Two electron system:

$$I(Q) = 2r_0^2 [1 + \cos(Qr)]$$


$$\mathbf{Q} \parallel \mathbf{r}$$

orientational average



“smeared”, no more “point-like” particles

for many systems, e.g. molecules the orientation of  $\mathbf{r}$  will be random wrt  $\mathbf{k}$



Orientational averaging: assume one electron at  $r=0$ , a second at  $r$

$$A(Q) = f_1 + f_2 \exp(iQr)$$

$$I(Q) = f_1^2 + f_2^2 + f_1 f_2 \exp(iQr) + f_1 f_2 \exp(-iQr)$$

orientational averaging:  $\langle \exp(iQR) \rangle = \langle \exp(-iQr) \rangle$

$$\langle I(Q) \rangle = f_1^2 + f_2^2 + 2f_1 f_2 \langle \exp(iQr) \rangle$$

$$\langle \exp(iQr) \rangle = \frac{\int \exp(iQrcos\theta) \sin\theta d\theta d\Phi}{\int \sin\theta d\theta d\Phi} \xrightarrow{4\pi}$$

$$\begin{aligned} & \int \exp(iQrcos\theta) \sin\theta d\theta d\Phi \\ &= 2\pi \int \exp(iQrcos\theta) \sin\theta d\theta \\ &= 2\pi (-1/iQr) \int_{iQr}^{-iQr} \exp(x) dx \\ &= 4\pi \sin(Qr)/Qr \end{aligned}$$

$$\langle I(Q) \rangle = f_1^2 + f_2^2 + 4\pi f_1 f_2 \sin(Qr)/Qr$$

see figure 4.2 b

if the position of the electrons distributed or smeared: see Figure 4.2c

# Scattering from an atom:

scattering amplitude of an atom  $\equiv$  atomic form factor  $f_0(Q)$  [in units of  $r_0$ ]

$\rho(r)$ : electronic number density  $\equiv$  charge density

$$f_0(Q) = \int \rho(r) \exp(iQr) dr$$

$$= \begin{cases} Z & Q \rightarrow 0 \\ 0 & Q \rightarrow \infty \end{cases}$$

note: atomic form factor is FT of electronic charge distribution

$f_0(Q/4\pi)$  tabulated:

$$f_0(Q/4\pi) = \sum_{j=1}^4 a_j \exp -b_j(Q/4\pi)^2 + c$$

	$a_1$	$b_1$	$a_2$	$b_2$	$a_3$	$b_3$	$a_4$	$b_4$	$c$
C	2.3100	20.8439	1.0200	10.2075	1.5886	0.5687	0.8650	51.6512	0.2156
O	3.0485	13.2771	2.2868	5.7011	1.5463	0.3239	0.8670	32.9089	0.2508
F	3.5392	10.2825	2.6412	4.2944	1.5170	0.2615	1.0243	26.1476	0.2776
Si	6.2915	2.4386	3.0353	32.333	1.9891	0.6785	1.5410	81.6937	1.1407
Cu	13.338	3.5828	7.1676	0.2470	5.6158	11.3966	1.6735	64.820	1.5910
Ge	16.0816	2.8509	6.3747	0.2516	3.7068	11.4468	3.683	54.7625	2.1313
Mo	3.7025	0.2772	17.236	1.0958	12.8876	11.004	3.7429	61.6584	4.3875

table 4.1: J. Als-Nielsen & D. McMorrow

note:

$$f = f_0(Q) + f' + f''$$


corrections  $f'$  and  $f''$  arise from the fact that the electrons are bound in the atom

# Scattering from a molecule:

$$F^{\text{mol}}(Q) = \sum r_j f_j(Q) \exp(iQr_j)$$

example: CF4:

assume OA=OB=OC=OD=1; z=OO'=cos(u) =1/3




$$Qr_j = Qr_j \cos(u) = (1/3)Qr_j$$

Assume:  $Q \parallel \text{C-F bond}$

$$\begin{aligned} F^{\text{mol}} &= f^C(Q) + f^F(Q) [\exp(iQR) + 3\exp(iQr_j)] \\ &= f^C(Q) + f^F(Q) [3\exp(-/+iQR/3) + \exp(+/-iQR)] \end{aligned}$$

- CF4
- - . - CF4  $Q \text{ not } \parallel \text{C-F}$
- - - - molybdenum  
(also 42 electrons)




(F-F bond)

# Scattering from a crystal

$$A(\mathbf{Q}) = -r_0 \sum r_j \exp(i \mathbf{Q} \mathbf{r}_j)$$

an extension to crystalline matter is simplified since there is translational symmetry.

crystalline matter:  $\mathbf{r}' = \mathbf{R}_n + \mathbf{r}_j$



$$A(\mathbf{Q}) = -r_0 \sum_{\mathbf{R}_n} \exp(i \mathbf{Q} \mathbf{R}_n) \sum r_j \exp(i \mathbf{Q} \mathbf{r}_j)$$

---

---

lattice sum

unit cell structure factor

## Crystallography:

determine electron density within unit cell

Note: one does measure  $I(Q) = A(Q) A^*(Q)$  and is thus not sensitive to phase shifts

# Scattering from atoms on a crystal lattice

concept: build up crystal from lattice+basis

$$2\text{-D lattice: } \mathbf{R}_n = n_1 \mathbf{a}_1 + n_2 \mathbf{a}_2$$

unit cell: primitive or (non-) primitive

(primitive  $\equiv$  area or volume minimized)

$$3\text{-D lattice: } \mathbf{R}_n = n_1 \mathbf{a}_1 + n_2 \mathbf{a}_2 + n_3 \mathbf{a}_3$$

There are: 5 distinct lattices in 2-D



14 distinct lattices in 3-D

called the Bravais lattices


add basis to yield:

32 possible point groups

230 possible symmetry groups




(d) Crystal=lattice+basis



# Lattice planes and Miller indices

The plane closest to the origin that has intercepts  $a_1/h$ ,  $a_2/k$ ,  $a_3/l$  is called the  $(h,k,l)$  plane



Rhombohedral,  $a = b = c$ ,  $\alpha = \beta = \gamma$ :

$$\frac{1}{d_{hkl}^2} = \frac{(h^2 + k^2 + l^2) \sin^2 \alpha + 2(hk + kl + lh)(\cos^2 \alpha - \cos \alpha)}{a^2(1 + 2 \cos^3 \alpha - 3 \cos^2 \alpha)}.$$

Hexagonal,  $a = b$ ,  $\alpha = \beta = 90^\circ$ ,  $\gamma = 120^\circ$ :

$$\frac{1}{d_{hkl}^2} = \frac{4}{3} \left( \frac{h^2 + hk + k^2}{a^2} \right) + \frac{l^2}{c^2}.$$

Monoclinic,  $\alpha = \gamma = 90^\circ$ :

$$\frac{1}{d_{hkl}^2} = \frac{1}{\sin^2 \beta} \left( \frac{h^2}{a^2} + \frac{k^2 \sin^2 \beta}{b^2} + \frac{l^2}{c^2} - \frac{2hl \cos \beta}{ac} \right).$$

Orthorhombic,  $\alpha = \beta = \gamma = 90^\circ$ :

$$\frac{1}{d_{hkl}^2} = \frac{h^2}{a^2} + \frac{k^2}{b^2} + \frac{l^2}{c^2}.$$

Tetragonal,  $a = b$ ,  $\alpha = \beta = \gamma = 90^\circ$ :

$$\frac{1}{d_{hkl}^2} = \frac{h^2 + k^2}{a^2} + \frac{l^2}{c^2}.$$


Cubic,  $a = b = c$ ,  $\alpha = \beta = \gamma = 90^\circ$ :

$$\frac{1}{d_{hkl}^2} = \frac{h^2 + k^2 + l^2}{a^2}.$$

**spacing formulae; B.E. Warren p.21**

# Laue conditions and reciprocal space

position of an atom in lattice and cell:



$$F_{\text{crystal}}(Q) = \sum_{r_j} F_j^{\text{mol}}(Q) \exp(iQr_j) \bullet \sum_{R_n} \exp(iQR_n)$$

---

unit cell structure factor

---

lattice sum

## Evaluation of the lattice sum:

$$\sum_{\mathbf{R}_n} \exp(i \mathbf{Q} \cdot \mathbf{R}_n) \quad (i)$$

number of terms is enormous

(i) is of order unity (many  $\exp(i\Phi)$  terms)

except for:

$$\mathbf{Q} \cdot \mathbf{R}_n = 2\pi \times \text{integer} \quad (ii)$$

in that case the lattice sum becomes big.

Concept: construct an artificial lattice with  
lattice vectors  $a_1^*$ ,  $a_2^*$  and  $a_3^*$   
such that:

$$a_i \cdot a_j^* = 2\pi \delta_{ij} \quad (iia)$$

then any vector defined on the  
reciprocal lattice is of type:

$$G = ha_1^* + ka_2^* + la_3^* \quad (iii)$$

(iii): G satisfies (ii) [ $\mathbf{Q} \cdot \mathbf{R}_n \times \text{integer}$ ]

since

$$\mathbf{G} \cdot \mathbf{R}_n = 2\pi (hn_1 + kn_2 + ln_3)$$

only if

$Q = G$

(Laue condition)

so that the scattering amplitude will not  
vanish.

Thus: Q need to be a reciprocal  
lattice vector.

# Reciprocal Lattice

1-D: defined by (iia):  $a_i \bullet a_j^* = 2\pi \delta_{ij}$

2-D and 3-D:

$$a_1^* = (2\pi/v_c) a_2 \times a_3$$

$$a_2^* = (2\pi/v_c) a_3 \times a_1$$

$$a_3^* = (2\pi/v_c) a_1 \times a_2$$

with  $v_c = a_1 \bullet (a_2 \times a_3)$

note: in 2-D  $a_3$  is chosen to be a unit vector normal to the 2-D plane spanned by  $a_1$  and  $a_2$ .


Example: fcc lattice

$$a_1 = (a/2)(y'+z'), a_2 = (a/2)(z'+x'), a_3 = (a/2)(x'+y')$$

$$a_1^* = (4\pi/a) (y/2 + z/2 - x/2)$$


$$a_2^* = (4\pi/a) (z/2 + x/2 - y/2)$$

$$a_3^* = (4\pi/a) (x/2 + z/2 - y/2)$$



# Equivalence of the Laue and the Bragg conditions

## (a) Equivalence of Bragg and Laue



Bragg's law:  $\sin \theta = \lambda/2d$


The Laue condition requires  
 $Q = G$

Example:

$$\begin{aligned} Q &= (0,1) = 0 \bullet a_1^* + 1 \bullet a_2^* \\ &= (2\pi/d) \underline{x}_2 \end{aligned}$$

$$\sin \theta = (Q/2)/k = Q\lambda/4\pi$$

$$\sin \theta = (2\pi/d)\lambda / 4\pi = \lambda/2d \text{ qed}$$



# The Ewald shere

Visualisation of diffraction effects in reciprocal space (a).

Laue condition requires  $Q = G = ha_1^* + ka_2^*$

Design sphere with radius  $k$  pointing to origin (b).


If any reciprocal lattice point falls on the circle then the Laue conditions is fulfilled (c).

Intensity is observed if the detector is placed in the direction of  $k'$  (c).

A rotation about O corresponds to a rotation of the crystal.

Note: More than one reciprocal lattice point can lie on the sphere  $\equiv$  multiple scattering.

If the beam is not monochromatic the sphere adopts the corresponding width. In the white beam case all spots are ultimately detected.



# The unit cell structure factor

$$F_{\text{uc}}(Q) = \sum_{rj} F_j^{\text{mol}}(Q) \exp(iQr_j)$$

example: fcc lattice (*use conventional cubic unit cell*)

$$a_1 = a\underline{x}, a_2 = a\underline{y}, a_3 = a\underline{z}$$

$$r_1 = 0, r_2 = \frac{1}{2}a(\underline{y} + \underline{z}), r_3 = \frac{1}{2}a(\underline{z} + \underline{x}), r_4 = \frac{1}{2}a(\underline{x} + \underline{y})$$

$$G = ha_1^* + ka_2^* + la_3^*$$

$$a_1^* = 2\pi/v_c (a_2 \times a_3) = 2\pi/a^3 [a\underline{y} \times a\underline{z}] = 2\pi/a [\underline{y} \times \underline{z}] = 2\pi/a \underline{x}$$

$$a_2^* = 2\pi/v_c (a_3 \times a_1) = 2\pi/a^3 [a\underline{z} \times a\underline{x}] = 2\pi/a [\underline{z} \times \underline{x}] = 2\pi/a \underline{y}$$

$$a_3^* = 2\pi/v_c (a_1 \times a_2) = 2\pi/a^3 [a\underline{x} \times a\underline{y}] = 2\pi/a [\underline{x} \times \underline{y}] = 2\pi/a \underline{z}$$


$$v_c = a \mathbf{1} \bullet (a_2 \times a_3)$$

$$G \bullet r_1 = 2\pi/a (hx + ky + lz) \bullet 0 = 0$$

$$G \bullet r_2 = 2\pi/a (h\underline{x} + k\underline{y} + l\underline{z}) \bullet 1/2a(\underline{y} + \underline{z}) = \pi (k+l)$$

$$G \bullet r_3 = 2\pi/a (h\underline{x} + k\underline{y} + l\underline{z}) \bullet 1/2a(\underline{z} + \underline{x}) = \pi (h+l)$$

$$G \bullet r_4 = 2\pi/a (h\underline{x} + k\underline{y} + l\underline{z}) \bullet 1/2a(\underline{x} + \underline{y}) = \pi (h+k)$$



# The unit cell structure factor for a fcc lattice

$$F_{hkl}^{\text{fcc}}(Q) = \sum_{j=1-4} f(Q) \exp(iQr_j) = f(Q) [ \exp(iGr_1) + \dots \exp(iGr_4) ]$$

$$F_{hkl}^{\text{fcc}}(Q) = f(Q) [ 1 + \exp(i\pi(k+l)) + \exp(i\pi(h+l)) + \exp(i\pi(h+k)) ]$$

$$= f(Q) \cdot \begin{cases} 4 & \text{if } h,k,l \text{ are all even or odd} \\ 0 & \text{otherwise} \end{cases}$$


$$I_{hkl}^{\text{fcc}}(Q) = F(Q) \bullet F^*(Q)$$

Reflections:

100 forbidden

111 allowed

200 allowed



# Lattice sums

$$S_N(Q) = \sum_{R_N} \exp(iQR_N)$$

1-D:  $R_N = na$

$$S_N(Q) = \sum_{n=0}^{N-1} \exp(iQna)$$

$$|S_n(Q)| = \sin(N\pi Q)/\sin(\pi Q)$$

Als-Nielsen&McMorrow p.51

Laue condition almost fulfilled:

$$Q = (h+\xi) a^*$$

$$S_N(\xi) = \exp(i \pi \xi (N-1)) \cdot [\sin(N\pi\xi) / \sin(\pi\xi)]$$

$$|S_N(\xi)| = \xrightarrow{N \text{ large}, \xi \rightarrow 0} N$$

width:

$$|S_N(\xi=1/2N)| \approx (2/\pi)N \\ \approx \frac{1}{2} (\text{peak height})$$

$$\text{FWHM} \sim 1/N$$

N large:

$$|S_N(\xi)| \longrightarrow \delta(\xi)$$

$$|S_N(Q)| \longrightarrow a^* \delta(Q - G_h)$$

$$|S_N(Q)|^2 \longrightarrow N a^* \delta(Q - G_h)$$

# Lattice sums (2D & 3D)

$$S_N(Q) = \sum_{RN} \exp(iQR_N)$$

2-D:

$$|S_N(\xi_1, \xi_2)| \longrightarrow N_1 N_2 \delta(\xi_1) \delta(\xi_2)$$

for large N1, N2:

$$|S_N(Q)|^2 \longrightarrow (N_1 a_1^*) (N_2 a_2^*) \delta(Q-G) = NA^* \delta(Q-G)$$

A\* area of unit cell in reciprocal space

3-D:

$$|S_N(Q)|^2 \longrightarrow N v_c^* \delta(Q-G)$$

with  $G=ha_1^*+ka_2^*+la_3^*$ , N: number of unit cells,  $v_c^*$  volume unit cell in reciprocal space

# Further topics

Quasiperiodic lattices

Crystal truncation rods

Lattice vibrations, Debye-Waller factor, TDS

Lorentz factor

Diffraction from a fiber

2-D crystallography