## Excercises "Methoden Moderner Röntgenphysik II"

Date, Room and Time: 13. April 2010, Seminarraum 3, 16:00-17:30

## — Crystal Truncation Rods —

In Born-Approximation, the x-ray amplitude scattered by a crystal is given by

$$A(\vec{q}) \propto S(\vec{q}) \sum_{n_1 = -M_1}^{N_1 - 1} \sum_{n_2 = -M_2}^{N_2 - 1} \sum_{n_3 = -M_3}^{N_3 - 1} \exp(i\vec{q} \cdot [n_1 \vec{a} + n_2 \vec{b} + n_3 \vec{c}])$$

with the structure factor  $S(\vec{q})$  the wave vector transfer  $\vec{q}$  and the unit cell vectors  $\vec{a}, \vec{b}, \vec{c}$ . At infinity crystals all  $M_k$  and  $N_k$  are equal to infinity. At semi-finite crystals with a surface e.g. in the  $\vec{a}, \vec{b}$ -plane  $M_3$  is a finite number e.g.  $M_3 = 0$ . For a cubic crystal and a variation of  $\vec{q}$  along  $\vec{c}$  (the z – axis) the  $\vec{a}$  - and  $\vec{b}$  -directions can be neglected. The problem becomes 1-dimensional.

1) Calculate the scattered intensity  $I(q_z)$  of the Crystal Truncation Rod CTR for the 1-dimensional semi-infinity crystal with the interface at  $M_3 = 0$  and finite but very large  $N_3$ .

Use 
$$I(q_z) = |A(q_z)|^2$$
 and 
$$\sum_{n=0}^{N-1} x^n = \frac{x^{N/2} - x^{-N/2}}{x^{1/2} - x^{-1/2}} \cdot x^{\frac{N-1}{2}}$$

2) Make a sketch of  $I(q_z)$  and explain the problems for  $N_3 \to \infty$ .

Introducing an absorption depth for the x-rays makes the calculations more realistic. The intensity of the x-rays in the depth z in an absorbing material follows  $I(z) = I_0 \exp(-z/\Lambda)$  with the absorption length  $\Lambda$ .

- 3) Modify the scattered amplitude  $A(q_z)$  of the Crystal Truncation Rod CTR for the 1-dimensional semi-infinity crystal with the interface at  $M_3=0$  introducing the absorption effects to each of the components of the sum.
- 4) Using  $N_3 \mid \vec{c} \mid >> \Lambda$  and  $\mid \vec{c} \mid << \Lambda$  (which is both usually the case) calculate the scattered intensity  $I(q_z)$  of the Crystal Truncation Rod CTR for the 1-dimensional semi-infinity crystal with the interface at  $M_3 = 0$  and finite but very large  $N_3$  including absorption. Make a sketch and compare with the sketch of 2).