

# Biology I: Structural Biology and Macromolecular Crystallography (MX)

Thomas R. Schneider, EMBL Hamburg

18/6/2013

[thomas.schneider at embl-hamburg.de](mailto:thomas.schneider@embl-hamburg.de)


# EMBL

- Funded 1974
- Inter-governmental organization
- 20 member states
- One associate state
- Missions
  - Basic Research in Molecular Biology
  - Technology and Instrumentation
  - Facilities and Services
  - Teaching and Training
  - Technology Transfer
- 5 sites
- 1400 employees from 60 nations
- Annual Budget ca. 85 M€



# Biology I to IV

- Structural Biology and Macromolecular Crystallography (MX)
- MX – The method
- MX – Collection and processing of diffraction data
- MX – Building models and the future



# Today

- What is a protein?
- The central dogma of molecular biology
- Milestones in macromolecular crystallography
  - The first structures
  - Recombinant production of proteins
  - Use of synchrotron radiation
  - Detector technologies
  - The protein data bank
  - Cryogenic sample cooling
- Crystallographic Workflow

# Look at Abl kinase

- 2148 atoms per molecule, 275 amino acids
- A protein consists of chained amino acids (20 types)
- Backbone vs. Sidechain
- alpha-helices
- beta-sheets
- Schematic 'cartoon' representations of proteins
- Surface representation of proteins (with properties mapped)
- Interactions with ligands can be studied
- Gleevec / Imatinib
  
- Folding (Stretch out 275 amino acids ->  $3.8\text{ cm} \times 275 = 10.41\text{ m}$ , compare to 53 Å / cm)

Downloads:


molecule viewer: <http://pymol.org/educational/>

pdb-model: <http://www.rcsb.org/pdb/explore/explore.do?structureId=1FPU>

viewer script: see course website


# A protein molecule

- Looking at a collection of >2000 atoms can be rather confusing:



# Polypeptide chain

- Proteins are heteropolymers of amino-acids:




# Protein Backbone

- Looking at the backbone only allows to follow the polypeptide chain across the protein



# Protein conformation

- alpha helices and beta sheets contain repeating pattern / hydrogen bonds:




# Protein conformation

- Schematic representation of helices and sheets illustrates the 'fold'




# Protein surfaces

- What other molecules 'see' is a molecular surface.
- Active sites are often found in cavities




# Active sites

- Knowledge of the active site geometry allows to design ligands binding to the active sites thus acting as drugs.
- Here: Gleevec binding Abl-Kinase




# Crystal structures can be used to understand the action of drugs



## The drug


Mortality rate reduced by 80% for patients where interferon did not work



Crystal structure of the drug bound to its target Abl-Kinase  
red shows resistant mutation

# The Central Dogma of Molecular Biology


- 'DNA makes RNA makes protein' (Nirenberg 2010)



<http://library.thinkquest.org/C0122429/intro/genetics.htm>

# Crystal Structures support Basic Biology

- RNA Polymerase II – Structural Basis of Transcription
- 12 proteins
- ~30000 atoms
- Nobel Prize 2006 to Roger Kornberg




<http://www.youtube.com/watch?v=6QMPU9nuQso>

<http://www.lmb.uni-muenchen.de/cramer/pr-materials/index.htm>


# Crystal Structures support Basic Biology

- Ribosome – Structural Basis of Translation
- 3 stretches of RNA, 52 proteins
- ~37000 RNA atoms,  
~22000 protein atoms
- Nobel Price 2009 to Steitz,  
Yonath, Ramakrishnan



# Some milestones

# DNA (1953)



Watson & Crick (1953) Nature 172:137


[http://www.nobelprize.org/nobel\\_prizes/medicine/laureates/1962/](http://www.nobelprize.org/nobel_prizes/medicine/laureates/1962/)

<http://commons.wikimedia.org/wiki/>

[File:DNA\\_Structure%2BKey%2BLlabelled.pn\\_NoBB.png](#)

# Myoglobin (1960)

- Started in 1954
- Structure at 2 Å resolution published in 1960
- Key-step: choose sperm-whale as source



KENDREW, J. C. et al. (1960) Nature, 185, 422–427.

[http://www.nobelprize.org/nobel\\_prizes/chemistry/laureates/1962/kendrew-lecture.html](http://www.nobelprize.org/nobel_prizes/chemistry/laureates/1962/kendrew-lecture.html)

pdb: 1MBN



# Crystals

- Are difficult to grow
- Are difficult to reproduce
- Are often very small (microns)
- Are often inhomogeneous
- Are mechanically fragile
- Are radiation sensitive




Here: Cueva de los Cristales,  
Chihuahua region, Mexico




# Finding Conditions



# Small Volume HTP Crystallization



96 x 50-200 nl



96 x 10 nl

# High-Throughput Crystallization at EMBL Hamburg


- Miniaturization
  - 200nl/exp for vapour diffusion
  - 10nl/exp for counter diffusion
- Reference:
  - J. Müller-Dieckmann (2006)  
Acta Cryst.  
D64:1146-1152.




Jochen Müller-Dieckmann  
Xandra Kreplin


# Small Crystals

- Crystal dimensions of 1-10 micron are not rare.
- Often these crystals are of high quality
- Small and parallel beams needed.




Coulibaly et al. The molecular organization of cypovirus polyhedra. Nature (2007) 446: 97-101






# Amyloid fibrils





Nelson et al. Structure of the cross-beta spine of amyloid-like fibrils.  
Nature (2005) 435: 773-8



# Recombinant production of proteins

- Insert DNA from other organism into bacteria (Lobban?, 1972) using enzymes as tools for the manipulation of DNA
- 1982 synthetic human insulin (Genentech & Eli Lilly) entered the market for diabetes therapy.
- Recombinant production (and 'overexpression') of protein molecules is crucial for macromolecular crystallography as large amounts of material are needed to produce crystals.



# Use of synchrotron radiation in Biology

434

NATURE VOL. 230 APRIL 16 1971

NATURE VOL. 230 APRIL 16 1971

## Synchrotron Radiation as a Source for X-ray Diffraction

G. ROSENBAUM &amp; K. C. HOLMES

Max-Planck-Institut für Medizinische Forschung, Heidelberg

J. WITZ

Laboratoire des Virus des Plantes, Institut de Botanique de la Faculté des Sciences de Strasbourg, Strasbourg

Some preliminary results have been obtained with synchrotron radiation from the 7.5 GeV electron synchrotron Deutsches Elektronen - Synchrotron (DESY) in Hamburg as a source for X-ray diffraction.

When an electron is accelerated it emits radiation. At the very high energies used in DESY, the emitted radiation is confined to a narrow cone about the instantaneous direction of motion of the electron. Thus the synchrotron radiation tangentially. Synchrotron radiation is polychromatic, with a peak in the X-ray region for an electron energy of 7.5 GeV (see ref. 1 for the original theoretical description and refs. 2-4 for experimental details).

The DESY synchrotron uses bursts of 50 pulses and each 10 ms pulse contains  $6 \times 10^{12}$  electrons (10 mA average beam current). The injection energy is relatively low and the electrons are accelerated up to 7.5 GeV in the 10 ms.

Most of the X-radiation is emitted during the last 3 ms of each pulse; little radiation is produced at the lower electron energies, and so the time averaged intensity at 1.5 Å is about 30% of the peak value.

Table 1 Data for Quartz Monochromator in Synchrotron Radiation Beam

Synchrotron Electron beam diameter	7.5 GeV, 10 mA beam current approximately 4 mm (= effective X-ray source diameter)
Distance from the incident beam to monochromator	37 m (from synchrotron to monochromator approximately $10^4$ rad)
Polarization	85% at 1.5 Å in the eighth ring of the cycle, polarized in the plane of the synchrotron
Be-window	0.5 mm ( $10^{-2}$ cm $^2$ )
Crystal	quartz (001) at $\theta = 8^\circ 30'$ to the 1011 planes, dimensions $45 \times 13 \times 0.3$ mm $^3$
Bender	pivot: outer pair 40.5 mm inner pair 39.5 mm
Wavelength	1.5 Å
Wavelength spread	$4 \times 10^{-4}$ Å (at 1.5 Å)
Focus	1.5 m from crystal
Angular aperture of emitted beam	horizontal: 2 mrad (isotropic)
Monochromator line focus	horizontal: 1.5 mrad (at the eighth ring of the cycle)

Table 2 Biological Applications

Specimen	Elliott fine-focus X-ray tube*	DESY synchrotron with Berlmann point-focusing monochromator†
Single crystal	Standard collimator 0.5 mm diameter	
$a = 0.5$ mm	$d = 12.5$ cm	$D = 1$ m
$b = 0.5$ mm	$d = 0.7$ mm	$d = 120$ µm
$L = 7.5$ cm	$P = 10^2$ photons s $^{-1}$	$P = 4 \times 10^2$ photons s $^{-1}$
	$J = 2 \times 10^2$ photons s $^{-1}$ mm $^{-2}$	$J = 2.5 \times 10^{11}$ photons s $^{-1}$ mm $^{-2}$
Tobacco mosaic virus gel	Double-crystal focusing monochromator‡	
$a = 0.5$ mm	$d = 80$ µm	$D = 0.8$ m
$b = 1$ mm	$P = 10^2$ photons s $^{-1}$	$d = 100$ µm
$L = 32$ cm	$J = 2 \times 10^2$ photons s $^{-1}$ mm $^{-2}$	$P = 1 \times 10^2$ photons s $^{-1}$
Insect muscle	Double-crystal focusing monochromator‡	
$a = 3$ mm	$d = 200$ µm	$D = 1.1$ (1.0) m
$b = 0.3$ mm	$P = 1 \times 10^2$ photons s $^{-1}$	$d = 180$ (200) µm
$L = 40$ cm	$J = 3 \times 10^2$ photons s $^{-1}$ mm $^{-2}$	$P = 3 \times 10^2$ (2 $\times$ 10 $^2$ ) photons s $^{-1}$ mm $^{-2}$

\* Width of specimen;  $a$ , height of specimen;  $b$ , specimen film distance;  $d$ , anode specimen distance;  $D$ , focal length; that is, monochromator film distance;  $d$ , spot or focus diameter on film;  $P$ , X-ray power radiating the specimen; and  $J$ , flux density at the focus.

† Loaded with 40 kV, 50 mA into a 0.1  $\times$  2 mm $^2$  electron focus at the anode of the rotating anode, and 40 kV, 15 mA into a 0.14  $\times$  0.1 mm $^2$  electron focus at the cathode focus. This set is the most powerful Berlmann X-ray tube currently available.

‡ The setting of this Johann-type‡ monochromator is optimized for each type of specimen.

\* Conditions of the synchrotron are as in Table 1, computed for 1.5 Å radiation.

We have evaluated the spectral luminance (that is, the power in photons per second radiated per unit area, solid angle, and wavelength interval) of both the synchrotron and a fine-focus rotating anode X-ray tube (see Table 2). The values are  $2 \times 10^{11}$  (time averaged) and  $3 \times 10^{22}$  photons s $^{-1}$  sterad $^{-1}$  cm $^{-2}$  Å $^{-1}$ , respectively at 1.54 Å, showing clearly that the synchrotron is, relative to present X-ray tubes, a very bright source. The actual advantage to be gained in a diffraction experiment depends critically on the optical system necessary to focus and monochromate the radiation. Three types of focusing mono-




Fig. 3 Equatorial reflexions from dorsolongitudinal flight muscle of *Lethocerus maximus* recorded with: *a*, monochromated synchrotron radiation; electron energy 5 GeV, beam current 8 mA, exposure time 15 min, specimen film distance 40 cm; note the parasitic scattering on the left of the backstop arising from fluorescence from the monochromator holder; *b*, Elliott fine-focus rotating anode tube at 40 kV, 15 mA, exposure time 1 h, specimen film distance 36 cm. The strong line is the 20 reflexion ( $d = 231$  Å); the weak lines are the 21, 31 and 32 reflexions.

# Use of synchrotron radiation in MX

## Applications of synchrotron radiation to protein crystallography: Preliminary results

(x-ray diffraction/anomalous dispersion/rubredoxin/azurin/nerve growth factor/glutaminase-asparaginase)

JAMES C. PHILLIPS, ALEXANDER WLODAWER, MARGUERITE M. YEVITZ, AND KEITH O. HODGSON\*

Department of Chemistry and Stanford Synchrotron Radiation Project, Stanford University, Stanford, California 94305

Communicated by Richard H. Holm, October 23, 1975

**ABSTRACT** X-ray diffraction photographs of protein single crystals have been obtained using synchrotron radiation produced by an electron-positron storage ring. The diffracted intensities observed with this unconventional source are a factor of at least 60 greater than those obtained with a sealed x-ray tube using the same crystal and instrumental parameters. Diffraction data have been collected by the precession method to higher resolution and using smaller protein crystals than would have been possible with a conventional source. The crystal decay rate in the synchrotron beam for several proteins appears to be substantially less than that observed with Ni-filtered Cu radiation. The tunable nature of the source (which allows selective optimization of anomalous contributions to the scattering factors) and the low angular divergence of the beam make the source very useful for single crystal protein diffraction studies.

The use of synchrotron radiation as a source for single crystal x-ray diffraction studies has recently been the subject of considerable discussion and controversy. In contrast to con-

tation of synchrotron radiation produced by the SPEAR electron-positron storage ring at the Stanford Linear Accelerator Center, Stanford, Calif.

Synchrotron radiation is emitted tangentially to the instantaneous path of the charged particles as they are main-




FIG. 3. Cone-axis oscillation photographs of the same azurin crystal. Precession angle 6.0°, oscillation angle 30°. (left) Synchrotron source,  $E = 3.7$  GeV,  $I = 40$  mA, only electrons present,  $\lambda = 1.346$  Å, exposure time 10 min. (right) Philips fine-focus sealed Cu anode tube, operated at 40 kV, 30 mA, exposure time 6 hr, Ni filter.


# Detectors

Technology	When	Readout	Remark
X-ray film	1940	30 min	grain size < 10 $\mu\text{m}$
Image Plates	1990	1.5 min	pixel: $150 \times 150 \mu\text{m}^2 + \text{PSF}$
CCD detectors	1997	1 sec	pixel: $80 \times 80 \mu\text{m}^2 + \text{PSF}$
Pixel-Array Detectors (2010)	2010	3 msec	pixel: $173 \times 173 \mu\text{m}^2$ sharp
Pixel Array Detectors (2014)	2104	??	pixel: $50 \times 75 \mu\text{m}$


- PSF = Point spread function

# The Protein Data Bank

- Started as a grass-root movement in the 70's
- As of Tuesday Jun 11, 2013 17:00 PDT, **91359 structures** are in the protein data bank.
- Out of these 80569 were determined by X-ray crystallography
- 1976:13
- 1980:69
- 1990: 507
- 2000: 13596
- 2010: 70013
- A large fraction of structures determined today are determined by 'Molecular Replacement'



# Radiation Damage



300μm beam

Garman & Schneider (1997) J. Appl. Cryst. 30:211

Thomas R. Schneider | Meth. moderner Röntgenphysik II | 18/6/2013

# Data collection at 100 K

- Mounting protein-crystals in a free-standing film revolutionized the field.

Teng, T. Y. (1990)  
J. Appl. Cryst 23: 387-391






Fig. 2. The loop mounted on an oscillation camera used at CHESS: *A* cold nitrogen nozzle; *B* the loop with a frozen crystal; *C* cold-gas-stream reflector mounted on the goniometer head. The cold nitrogen nozzle (*A*) of a transfer line is 1/4 in in diameter which delivers a gas stream at from 80 to 230 K. Within a cone-shaped working volume of 65 mm<sup>3</sup>, the temperature gradient is less than 4 K, at an initial exit temperature of 85 K. Fog or ice formation around the nozzle, crystal and goniometer head is avoided by a coaxial warm and dry nitrogen stream that surrounds the cold stream, and by a built-in heater on the base of the cold-gas-stream reflector (*C*).




# HPGonioV: Sample environment



- 400 micron thick blade.
- Motorized adjustable distance to sample: 7-36 mm.



# Radiation Damage

- Data collection at 100 K significantly extends crystal lifetime




Garman & Schneider (1997) J. Appl. Cryst. 30:211

# Radiation damage at 100 K




- On modern synchrotron beamlines the lifetime of protein crystals at 100k is on the order of seconds to minutes.
- Going back to Room temperature, Owen et al. (2012) Acta Cryst. D68:810 have shown that 'in the first 200 msec of a room temperature experiment, one can outrun hydroxyl radicals.'



# Crystal Structure Determination

# Diffraction from a Crystal




# Inside a crystal



- A crystal consists of repeating units, the crystallographic unit cells.
- Each unit cells has 'the same' content, i.e. the same molecules in the same conformation and in the same orientation
- In real crystals, there is always some amount of 'disorder'


# Diffraction from a crystal



- When electromagnetic waves are interacting with a periodic structure, interference effects will occur.

# Diffracting planes

[http://en.wikipedia.org/wiki/X-ray\\_crystallography](http://en.wikipedia.org/wiki/X-ray_crystallography)



Nobel prize for physics 1914 to Max von Laue




Nobel prize for physics 1915 to William and Lawrence Bragg




Constructive interference occurs when Bragg's law is fulfilled:

$$2 d \sin\theta = n \lambda$$



# Diffraction Data

- The diffraction pattern changes when the crystal is rotated.
- By rotating the crystal, different Bragg-planes are brought into their diffracting position



[http://www-structmed.cimr.cam.ac.uk/Course/  
Basic\\_diffraction/data\\_animation.html](http://www-structmed.cimr.cam.ac.uk/Course/Basic_diffraction/data_animation.html)

# Diffraction and forests ...




[http://flickr.com/photos/  
rossogiallobianco/2486114038](http://flickr.com/photos/rossogiallobianco/2486114038)

Thomas R. Schneider | Meth. moderner Röntgenphysik II | 18/6/2013

# Indices and Structure factor amplitudes

- Every diffraction spot is marked by an index  $hkl$
- For every diffraction spot an Intensity  $I$  is measured.
- The result of the experiment is an indexed set of  $I$ 's
- Diffraction at low  $2\theta$  is stronger than at high  $2\theta$



# Diffraction data


<b>h</b>	<b>k</b>	<b>l</b>	<b>I</b>	<b>sig(I)</b>
0	20	35	4980.5	122.6
0	20	36	6906.6	216.6
0	20	37	8302.3	231.7
0	20	38	3209.5	89.3
0	20	39	459.6	22.1
0	20	40	1017.4	33.8
0	20	41	-5.6	18.3
0	20	42	33.8	15.6
0	20	43	4545.7	133.4
0	20	44	210.5	19.2
0	20	45	808.8	29.1

# Structure Factor Amplitudes

- For formal reasons, the measured I's are usually converted to 'Structure Factor Amplitudes' F by:

$$F = \text{sqrt}(I).$$

# Calculating Electron density



$$\rho_{xyz} = \sum_{hkl} |F_{hkl}| e^{-i\varphi_{hkl}} e^{-2\pi i(hx+ky+lz)}$$

Electron density at some point xyz in space


Sum over all Bragg reflections

**Structure Factor Amplitude** for each reflection hkl.  $F_{hkl} \approx \sqrt{I_{hkl}}$



**Structure Factor Phase** for each reflection hkl.

**Phase shift** depending on hkl and position in space

# Typical initial electron density map



# Interpretation of the electron density map



# A 'pdb'-file ([www.rcsb.org](http://www.rcsb.org))

HEADER TRANSFERASE  
TITLE STRUCTURAL BASIS FOR THE AUTO-INHIBITION OF C-ABL TYROSINE  
TITLE 2 KINASE

			X	Y	Z	B		
.								
.								
ATOM	1	N	ALA A 243	20.064	-2.529	43.315	1.00 51.64	N
ATOM	2	CA	ALA A 243	19.658	-1.370	42.459	1.00 51.36	C
ATOM	3	C	ALA A 243	20.832	-0.838	41.643	1.00 50.91	C
ATOM	4	O	ALA A 243	20.650	0.010	40.776	1.00 50.77	O
ATOM	5	CB	ALA A 243	19.092	-0.263	43.322	1.00 50.20	C
ATOM	6	N	MET A 244	22.030	-1.350	41.906	1.00 50.62	N
ATOM	7	CA	MET A 244	23.218	-0.888	41.195	1.00 50.73	C
ATOM	8	C	MET A 244	23.537	-1.709	39.950	1.00 50.43	C
ATOM	9	O	MET A 244	24.554	-1.488	39.298	1.00 50.33	O
ATOM	10	CB	MET A 244	24.420	-0.882	42.146	1.00 51.32	C
ATOM	11	CG	MET A 244	24.215	0.020	43.361	1.00 52.32	C
ATOM	12	SD	MET A 244	25.597	0.070	44.515	1.00 54.49	S
ATOM	13	CE	MET A 244	26.730	1.130	43.627	1.00 53.37	C
ATOM	14	N	ASP A 245	22.662	-2.653	39.620	1.00 49.87	N
ATOM	15	CA	ASP A 245	22.857	-3.505	38.446	1.00 49.28	C
ATOM	16	C	ASP A 245	22.115	-2.895	37.257	1.00 47.55	C